

# BETA-DECAY AND DELAYED NEUTRON EMISSION OF VERY NEUTRON-RICH NUCLEI \*

I. N. Borzov

*Institut d'Astronomie et d'Astrophysique, Université Libre de Bruxelles,  
CP226, Bvd. du Triomphe, Belgium*

*and*

*University of Leuven, B-3001 Leuven, Belgium*

The weak interaction rates of very neutron-rich nuclei are among the most poorly known ingredients needed for stellar r-process modeling. Recently the nuclei near the closed neutron shells at  $N=50,82,126$  have been intensively studied at the RIB facilities. Comparison with the experimental data shows that the self-consistent framework can provide a reasonably sound prediction of the weak rates in the regions far from stability.

We briefly describe the continuum QRPA approach in terms of the finite Fermi-system theory. The ground state properties are treated within the nuclear energy-density functional framework [1,2]. The  $\beta$ -decay strength functions are calculated taking into account the Gamow-Teller and first-forbidden transitions [3].

The impact of the  $T=1$  and  $T=0$  pairing on the  $\beta$ -decay and delayed neutron emission rates has been studied. A performance of existing global models for the nuclides near the r-process paths at  $N=50, 82, 126$  is critically analyzed and confronted with the recent RIB experiments in the regions of  $^{78}\text{Ni}$  [4],  $^{132}\text{Sn}$  [5] and "east" of  $^{208}\text{Pb}$  [6].

\* Supported by the PAI Program IAP 5/07 "Exotic Nuclei for Nuclear Physics and Astrophysics".

- [1] S. A. Fayans *et al.*, Nucl. Phys. **A676**, 49 (2000).
- [2] I. N. Borzov and S. Goriely, Phys. Rev. C **62**, 035501 (2000).
- [3] I. N. Borzov, Phys. Rev. C **67**, 025802 (2003).
- [4] S. Franchoo *et al.*, Phys. Rev. Lett. **81**, 3100 (1998).
- [5] J. Shergur *et al.*, Phys. Rev. C **65**, 034313 (2002).
- [6] H. De Witte, A. N. Andreev, I. N. Borzov *et al.*, Phys. Rev. C (in print) 2004.